

Guide to vaccines for children who were born prematurely

Most vaccines are given as an injection in the thigh or upper arm. Rotavirus vaccine is given as drops to be swallowed.

Age due	Diseases protected against		Vaccine given
Eight weeks old	Diphtheria, tetanus, pertussis (whooping cough), polio, Haemophilus influenzae type b (Hib) and hepatitis B		DTaP/IPV/Hib/HepB
	Meningococcal group B (MenB)		MenB
	Rotavirus gastroenteritis		Rotavirus ²
Twelve weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B		DTaP/IPV/Hib/HepB
	MenB		MenB
	Rotavirus		Rotavirus ²
Sixteen weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B		DTaP/IPV/Hib/HepB
	Pneumococcal (13 serotypes)		Pneumococcal conjugate vaccine (PCV)
One year old (on or after child's first birthday)	Pneumococcal		PCV booster
	Measles, mumps and rubella (German measles)		MMR ¹
	MenB		MenB booster
Eighteen months of age	Born before 1 July 2024 No appointment	Born on or after 1 July 2024 DTaP/IPV/Hib/HepB Measles, mumps, rubella	DTaP/IPV/Hib/HepB MMR ¹
Three years four months old or soon after	Born before 1 July 2024 Diphtheria, tetanus, pertussis and polio Measles, mumps, rubella	Born on or after 1 July 2024 Diphtheria, tetanus, pertussis and polio	dTaP/IPV MMR ¹

1. In the UK, we have 2 MMR vaccines which work very well. One of them contains porcine gelatine and the other one doesn't. If you would prefer to have the vaccine that does not contain porcine gelatine, talk to your practice nurse or GP. You can view the MMR vaccine Patient Information Leaflets at: Priorix: www.medicines.org.uk/emc/product/1159/pil. MMRVAXPRO: www.medicines.org.uk/emc/product/6307/p

2. Rotavirus vaccine should only be given after checking for SCID screening result.

Further information

A guide to vaccinations at one year of age contains more detailed information about the routine childhood immunisation programme including the types of vaccine. Ask your health visitor for a copy if you were not given one soon after the birth of your baby.

From birth onwards, children born prematurely should also continue to follow the normal schedule see the leaflets:

- **A guide to vaccinations for children aged 18 months**
- **Pre-school vaccinations: guide to vaccinations from 2 to 5 years**
- **Immunisations for young people**

You can also visit: www.nhs.uk/vaccinations

Each vaccine has a Patient Information Leaflet, you can view them at www.medicines.org.uk



Vaccines are offered at the right time for your baby when they most need to develop the protection. Having the vaccines on time, helps to give them the best protection from these infectious diseases.



© Crown copyright 2025
PB25EN001 1p 20K MAY 2025

First published August 2006

Updated July 2025

To order more free copies or any other vaccination information, visit:
www.healthpublications.gov.uk
or phone: 0300 123 1002 (8am to 6pm, Monday to Friday)



UK Health
Security
Agency



UPDATED SCHEDULE FOR 2025

A quick guide to childhood vaccinations for the parents of

premature babies



i mmunisation
the safest way to protect your child

‘The two public health interventions that have had the greatest impact on the world’s health are clean water and vaccines’

– World Health Organization

What is immunisation and why is it needed?

Immunisation is a way of protecting ourselves from serious diseases. There are some diseases that can kill children or cause lasting damage to their health. Immunisations are given to prepare your child’s immune system (their natural defence system) to fight off those diseases when your child comes into contact with them.

When should my baby be immunised?

It is important that your baby has their vaccinations at the right age (see the back cover). This will help to keep the risk of your baby catching a serious disease as low as possible. The risk of side effects from some vaccines may increase if you put them off.

My baby was born early. When should premature babies have their first immunisation?

Premature babies have a higher risk of infection. They should be immunised in line with the recommended schedule from 8 weeks after birth, no matter how premature they were. This may happen whilst your baby is in hospital, you will need to discuss this with your doctor.

What happens at the appointment?

The doctor or nurse will explain the immunisation process to you, and answer any questions you have. The vaccine is injected into the muscle of the thigh. If your baby was born very prematurely then they may still be in hospital when the first routine immunisation is due.

Are there any reasons why my baby should not be immunised?

There are very few reasons why babies cannot be immunised. The vaccines should not be given to babies who have had:

- a confirmed anaphylactic reaction (severe allergic reaction) to a previous dose of the vaccine
- a confirmed anaphylactic reaction to neomycin, streptomycin, or polymyxin B (antibiotics used in vaccines) or
- who are suspected to have SCID or have a positive SCID test result

If your baby’s immune system is ‘suppressed’ (because they are having treatment for a serious condition such as a transplant or cancer) or if you have taken certain medications whilst pregnant which suppressed your immunity, then your baby may not be able to have certain vaccines such as Rotarix for rotavirus. Your doctor or practice nurse should get advice from a specialist.

There are no other medical reasons why these vaccines should definitely not be given.

Dealing with common side effects

There may be redness, soreness or tenderness where the injection is given and a few babies may develop a mild fever. Make sure you keep your child cool by:

- giving them plenty of fluids
- giving infant liquid paracetamol – check the dose with your doctor
- making sure they don’t have too many layers of clothes or blankets on

If your child’s face feels hot to the touch, or if your child becomes ill, trust your instincts and ask your doctor for advice. Or call **NHS 111**.

For more information about side effects, see **A guide to vaccinations at one year of age**.

Watch out for meningitis and septicaemia (blood poisoning)

Hib, MenB and pneumococcal vaccines protect against the most important bacteria causing meningitis and septicaemia in children and young people. However, as these diseases can be caused by many other bacteria and viruses, it is important to know the signs and symptoms of these illnesses.

Early symptoms of meningitis include fever, being irritable and restless, vomiting and refusing food – symptoms that are also common with colds and flu. But a baby with meningitis or septicaemia can become seriously ill within hours.

Look out for **one or more** of the following symptoms:

- a high-pitched, moaning cry
- irritable when picked up
- drowsy, difficult to wake
- stiff with jerky movements (convulsions/fits)
- pale, blotchy skin or turning blue
- fever, with cold hands and feet
- red or purple spots that do not fade under pressure. (Do the glass test explained below). On dark skin, check inside the eyelids or roof of the mouth where the spots may be more visible.



Press the side of a clear glass against the rash and see if the rash fades and loses colour. If it doesn’t, contact your doctor immediately. On dark skin, check inside the eyelids or roof of the mouth where the spots may be more visible.

If your child becomes ill with one or more of the signs or symptoms described above, contact your doctor urgently.

If you are still worried after getting advice, trust your instincts and take your child to your nearest hospital with an emergency department.